

Hydrodynamic Equation for the Breakdown of the Quantum Hall Effect in a Uniform Current *

Hiroshi AKERA**

Department of Applied Physics, Hokkaido University, Sapporo 060-8628

(Received October 16, 2000)

The hydrodynamic equation for the spatial and temporal evolutions of the electron temperature T_e in the breakdown of the quantum Hall effect at even-integer filling factors in a uniform current density j_x is derived from the Boltzmann-type equation, which takes into account electron-electron and electron-phonon scattering processes. The derived equation has a drift term, which is proportional to j_x and the first spatial derivative of T_e . When applied to the spatial evolution of T_e in a sample with an abrupt change of the width along the current direction, the equation gives a distinct dependence on the current direction as well as a critical relaxation, in agreement with results of recent experiments.

KEYWORDS: integer quantum Hall effect, nonlinear transport, theory

The diagonal conductivity σ_{xx} is extremely small in the quantum Hall effect^{1,2)} (QHE) at small currents. When the current is increased up to a critical value, σ_{xx} increases by several orders of magnitude within a narrow range of current, and the QHE breaks down^{3,4,5)} (reviews are given in refs. 6, 7, 8). The hysteresis^{3,4,5)} with the change of the current as well as the variety of spatial^{9,10,11,12)} and temporal^{4,13)} behaviors observed in the breakdown of the QHE can be understood to be caused by a nonequilibrium phase transition. One of the most promising theories for describing the nonequilibrium phase transition is a hot-electron theory,^{3,14,15)} in which the change of σ_{xx} is ascribed to the change of the electron temperature T_e . In this research, the spatial evolution of σ_{xx} in the current direction is studied on the basis of the hot-electron theory.

The spatial evolutions of σ_{xx} have been studied experimentally^{9,10,11,12)} in a system with an abrupt change of the sample width at $x = 0$ (the x axis is taken along the current), which is designed to produce the change of the current density $\mathbf{j} = (j_x, 0)$ at $x = 0$. When the sample is designed so that $|j_x(x < 0)| < j_{c1} < j_{c2} < |j_x(x > 0)|$, with j_{c1} and j_{c2} the critical current densities (the system is bistable at $j_{c1} < |j_x| < j_{c2}$), the relaxation in $x > 0$ of σ_{xx} to the larger stable value and the development of the breakdown have been observed.^{9,10,12)} On the other hand, when $|j_x(x < 0)| > j_{c2} > j_{c1} > |j_x(x > 0)|$, the relaxation in $x > 0$ of σ_{xx} to the smaller stable value and the recovery of the QHE have been observed.¹¹⁾ A unique feature in this nonequilibrium phase transition is that the length scale for the relaxation of σ_{xx} depends strongly on the current direction: it is much larger for the electron flow in the $+x$ direction. It is only for this direction of the electron flow that the relaxation length in $x > 0$ diverges as $j_x(x > 0)$ approaches a critical value.^{10,11,12)} According to the hot-electron theory, this

current-direction dependence can be understood by the drift motion of hot electrons along the current.^{9,11)} In order to understand the experiments and other spatial and temporal evolutions in a unified manner, the general theoretical framework is highly desirable.

In this study, we derive the hydrodynamic equation describing the spatial and temporal evolution of T_e from the Boltzmann-type equation on the basis of the microscopic scattering processes proposed previously.¹⁶⁾ We restrict our derivation to even-integer filling factors $2N$ ($N = 1, 2, \dots$), for simplicity, and assume a uniform current distribution. We apply the obtained hydrodynamic equation to calculate the spatial relaxation of T_e in the current direction. Our calculation reproduces both the current direction dependence and the critical behavior of the relaxation length.

From a microscopic point of view, the potential in the plane of the two-dimensional electron system (2DES) fluctuates due to the presence of the layer with ionized donors,¹⁷⁾ and the average distance between a potential hill and a neighboring valley, l_{vh} , is $l_{vh} \sim 0.1 \mu\text{m} \gg \ell$ with ℓ the magnetic length. The screened potential has an amplitude approximately equal to the Landau level separation,^{18,19)} and electrons (holes) populate the $N + 1$ th (N th) Landau level, even with a filling factor of $2N$. Electron orbits in such potential consist of localized closed orbits around a potential hill (valley) and extended orbits at the center of the broadened Landau level. Wave packets constructed from extended states perform drift motions perpendicular to the electric field. In addition, an electron hops between a large hill orbit and a large valley orbit near an extended orbit. Such a hopping motion is given by an electron-electron scattering, in which one electron hops between a hill and a valley and, at the same time, another is excited or relaxed within a hill or a valley. Hopping motions along the electric field excite electrons, resulting in the energy gain of the 2DES. Electron-phonon scatterings, on the

* to be published in J. Phys. Soc. Jpn. **70** (2001) No.6.

** E-mail address: akera@eng.hokudai.ac.jp

other hand, relax the 2DES, resulting in the energy loss. In our previous study,¹⁶⁾ the energy gain and loss have been calculated for the higher T_e branch and the vanishing lattice temperature in homogeneous states. The theory is generalized in this study to include spatial variations of T_e in a macroscopic scale (much larger than l_{vh}) as well as an effect due to the drift motion of electrons in extended orbits (this effect is absent in homogeneous states).

In constructing the Boltzmann-type equation, the 2DES is divided into regions separated by an equipotential line of the fluctuating potential so that each region contains one hill or one valley. The distribution of electrons is described by the occupation probability $f_{\alpha ia}(t)$ of the a th state in the i th region at position $\mathbf{r}_i$ in level α , where α is a combination of the Landau index and the spin and takes one of $N \uparrow, N \downarrow, N+1 \uparrow$, or $N+1 \downarrow$. The Boltzmann-type equation is then given to be

$$\frac{\partial f_{\alpha ia}}{\partial t} = \left(\frac{\partial f_{\alpha ia}}{\partial t} \right)_{\text{drift}} + \left(\frac{\partial f_{\alpha ia}}{\partial t} \right)_{\text{coll}}, \quad (1)$$

consisting of the drift term due to extended states and the collision term due to electron-phonon and electron-electron interactions. In electron-phonon scatterings, we have only included transitions $\alpha ia \rightarrow \alpha ib$ within the same region (i) within the same Landau level (α), since their rates are much larger than others due to larger overlaps of wave functions.¹⁶⁾ In electron-electron scatterings, we have only included transitions $\alpha ia \rightarrow \alpha jb, \beta kc \rightarrow \beta ld$ within the same Landau level (α, β). In the following we consider two classes of dominant electron-electron scatterings: (e1) $i = j$ and $k = l$, (e2) $i = j$, and k and l are the nearest neighbors.

The local equilibrium is assumed in the scale of l_{vh} , since the inelastic scattering length due to electron-electron scatterings is estimated to be $\sim \ell \ll l_{\text{vh}}$. Introducing the electron temperature $T_e(\mathbf{r}_i, t)$ and the chemical potential $\mu(\mathbf{r}_i, t)$ of the i th region, the occupation probability is given by

$$f_{\alpha ia}(t) = f(\varepsilon_{\alpha ia}; \mu(\mathbf{r}_i, t), T_e(\mathbf{r}_i, t)), \quad (2)$$

where $f(\varepsilon; \mu, T_e) = 1/\{\exp[(\varepsilon - \mu)/k_B T_e] + 1\}$, and $\varepsilon_{\alpha ia}$ is the energy of state αia .

A state in the local equilibrium is specified by the electron temperature $T_e(\mathbf{r}, t)$, the chemical potential $\mu(\mathbf{r}, t)$, and the electric field $\mathbf{E}(\mathbf{r}, t)$ (these are assumed to be slowly varying in the scale of l_{vh}). The hydrodynamic equations for these variables are coupled in general. Here we assume that the current density is uniform, neglecting deviations near the sample edges. Although the current distribution in the Hall bar geometry is not uniform in the QHE,²⁰⁾ it is suggested to be uniform in the breakdown region by the observations in many samples^{21, 22)} that the critical current is proportional to the sample width. $\mathbf{E}$ is then given by $\mathbf{E} = (\rho_{xx} j_x, \rho_{yx} j_x)$, with a constant current density $\mathbf{j} = (j_x, 0)$. Since ρ_{xx} is smaller than ρ_{yx} by, at least, one order of magnitude and ρ_{yx} is quantized even in the breakdown regime, $\mathbf{E}$ can be approximated by $\mathbf{E} = (0, -j_x h/2e^2 N)$. The chemical potential is then at the middle point between the N th

and the $N+1$ th Landau levels and $\nabla \mu = e\mathbf{E}$. Owing to the assumed uniformity of the current, the equation for $T_e(x, t)$ is decoupled from those for $\mathbf{E}$ and μ .

To obtain the equation for $T_e(x, t)$, we introduce the energy density $U(x, t)$ defined by

$$U(x, t) = \frac{1}{S} \sum_{\alpha, i, a} (\varepsilon_{\alpha ia} - \mu_i) f_{\alpha ia}(t), \quad (3)$$

with $\mu_i = \mu(\mathbf{r}_i)$. An average is taken over regions within area $S (\gg l_{\text{vh}}^2)$ in which the variation of T_e is negligible. The time derivative of $U(x, t)$ is given in terms of the electronic specific heat $C_e = \partial U / \partial T_e$ as

$$\frac{\partial U}{\partial t} = C_e \frac{\partial T_e}{\partial t}, \quad (4)$$

and is also given using eq. (1) as

$$\frac{\partial U}{\partial t} = \left(\frac{\partial U}{\partial t} \right)_{\text{drift}} + \left(\frac{\partial U}{\partial t} \right)_{\text{coll}}^{\text{p}} + \left(\frac{\partial U}{\partial t} \right)_{\text{coll}}^{\text{e1}} + \left(\frac{\partial U}{\partial t} \right)_{\text{coll}}^{\text{e2}}. \quad (5)$$

The three collision terms are the electron-phonon term, and the electron-electron terms (e1) and (e2), respectively.

The drift term is given by

$$\left(\frac{\partial U}{\partial t} \right)_{\text{drift}} = -\frac{\partial j_{Ux}^{\text{d}}}{\partial x}, \quad (6)$$

with the energy flux $\mathbf{j}_U^{\text{d}} = (j_{Ux}^{\text{d}}, 0)$:

$$j_{Ux}^{\text{d}} = 2\hbar\omega_c f_{\text{ex}} j_{nx}^{\text{d1}}, \quad (7)$$

where $j_{nx}^{\text{d1}} = eE_y/h$ is the electron flow averaged within $S (\gg l_{\text{vh}}^2)$ at the filling factor one, $f_{\text{ex}} = 1/[\exp(\hbar\omega_c/2k_B T_e) + 1]$ is the occupation probability of extended states, and ω_c is the cyclotron frequency. The electron-phonon term is given by

$$\begin{aligned} \left(\frac{\partial U}{\partial t} \right)_{\text{coll}}^{\text{p}} &= \frac{1}{S} \sum_{\alpha, i, a, b} (\varepsilon_{\alpha ib} - \varepsilon_{\alpha ia}) f_{\alpha ia} (1 - f_{\alpha ib}) W_{\alpha ia \rightarrow ib}^{\text{p}} \\ &= -P_L, \end{aligned} \quad (8)$$

where $W_{\alpha ia \rightarrow ib}^{\text{p}}$ is the transition rate and P_L is the energy loss per unit area per unit time. When $|\varepsilon_{\alpha ia} - \varepsilon_{\alpha ib}| \ll k_B T_e$ (this inequality holds approximately in the higher T_e branch ($T_e \geq T_{c1}$) since $|\varepsilon_{\alpha ia} - \varepsilon_{\alpha ib}| \sim 0.1\hbar\omega_c$ and $k_B T_{c1} = 0.324\hbar\omega_c$ ¹⁶⁾), we have $f_{\alpha ia}(1 - f_{\alpha ib}) \approx f_{\alpha ia}(1 - f_{\alpha ia}) = -k_B T_e f'(\varepsilon_{\alpha ia})$ and obtain

$$P_L = C_L \tilde{T}_e, \quad (9)$$

with $\tilde{T}_e = k_B T_e / \hbar\omega_c$ and

$$C_L = \frac{\hbar\omega_c}{S} \sum_{\alpha, i, a, b} [-f'(\varepsilon_{\alpha ia})](\varepsilon_{\alpha ia} - \varepsilon_{\alpha ib}) W_{\alpha ia \rightarrow ib}^{\text{p}}. \quad (10)$$

The electron-electron term (e1) is given, when $\partial T_e / \partial x \ll T_e / l_{\text{vh}}$, by

$$\left(\frac{\partial U}{\partial t} \right)_{\text{coll}}^{\text{e1}} = \frac{\partial}{\partial x} \kappa_0 \frac{\partial T_e}{\partial x}, \quad (11)$$

where κ_0 is the thermal conductivity due to (e1) pro-

cesses. When $|\varepsilon_{\alpha ia} - \varepsilon_{\alpha ib}| \ll k_B T_e$, κ_0 is given by

$$\kappa_0 = \frac{k_B}{8S} \sum_{\alpha, \beta, i, k, a, b, c, d} W_{\beta kc \rightarrow kd}^{\alpha ia \rightarrow ib} [-f'(\varepsilon_{\alpha ia})] [-f'(\varepsilon_{\beta kc})] \Delta \varepsilon^2 \Delta r^2, \quad (12)$$

with $\Delta \varepsilon = \varepsilon_{\alpha ia} - \varepsilon_{\alpha ib}$, $\Delta r = |\mathbf{r}_i - \mathbf{r}_k|$, and $W_{\beta kc \rightarrow kd}^{\alpha ia \rightarrow ib}$ the transition rate. The electron-electron term (e2) is given by

$$\left(\frac{\partial U}{\partial t} \right)_{\text{coll}}^{\text{e2}} = P_G - \frac{\partial j_{Ux}^h}{\partial x}, \quad (13)$$

where $P_G = (-e)\mathbf{E} \cdot \mathbf{j}_n^h$ is the energy gain per unit area per unit time, and $\mathbf{j}_n^h$ and $\mathbf{j}_U^h = (j_{Ux}^h, 0)$ are the electron flux and the energy flux due to hopping processes, respectively. When $|\varepsilon_{\alpha ia} - \varepsilon_{\alpha ib}| \ll k_B T_e$ and $eEl_{\text{vh}} \ll k_B T_e$ (the second inequality holds at $E \sim E_{c1}$ since $E_{c1} = 20\text{V/cm}$ ($B = 5\text{ T}$)¹⁶ and $eE_{c1}l_{\text{vh}} = 0.03\hbar\omega_c$), we obtain

$$P_G = C_G f_{\text{ex}}(1 - f_{\text{ex}}) \tilde{E}^2, \quad (14)$$

where $\tilde{E} = eEl/\hbar\omega_c$ (ℓ the magnetic length) and

$$C_G = \frac{(\hbar\omega_c)^2 l_{\text{vh}}^2}{4\ell^2 S} \sum_{\alpha, \beta, i, k, l, a, b, c, d} W_{\beta kc \rightarrow ld}^{\alpha ia \rightarrow ib} [-f'(\varepsilon_{\alpha ia})]. \quad (15)$$

Similarly we obtain

$$j_{Ux}^h = -C_G f_{\text{ex}}(1 - f_{\text{ex}}) \frac{\ell^2}{4T_e} \frac{\partial T_e}{\partial x}. \quad (16)$$

Finally we obtain the equation for T_e :

$$C_e \frac{\partial T_e}{\partial t} = \frac{\partial}{\partial x} \kappa(T_e) \frac{\partial T_e}{\partial x} - \frac{\partial j_{Ux}^d}{\partial x} + P_G - P_L, \quad (17)$$

where the total thermal conductivity κ is

$$\kappa(T_e) = \kappa_0 + C_G f_{\text{ex}}(1 - f_{\text{ex}}) \frac{\ell^2}{4T_e}. \quad (18)$$

When a constant density of states is assumed, C_L , C_G , and κ_0 are approximately independent of T_e . Estimates of C_L and C_G are given in our previous paper.¹⁶ In estimating C_L , the deformation potential scattering by acoustic phonons and the vanishing lattice temperature were assumed; in estimating C_G , the summation over k and l was limited to the case where either k or l coincides with i . Calculated values of the critical electric field $E_{c1} \propto (C_L/C_G)^{1/2}$ and the energy dissipation $\sigma_{xx} E_{c1}^2 \propto C_L$ were in agreement with the observed values in their orders of magnitude. Following the calculation of C_G , here, we calculate κ_0 , in which we limit the summation over k to the nearest neighbors of i since the contribution to κ_0 of larger Δr decreases as Δr^{-3} . The estimation shows that κ_0 is only 5 percent of the total thermal conductivity, κ . The T_e dependence of κ is neglected in the following, since $\partial \kappa / \partial T_e = 0$ at $E = E_{c1}$ and $T_e = T_{c1}$.

Gurevich and Mints¹⁴) introduced an equation for T_e in their study of spatial and temporal variations of T_e in the bistable region. In their equation they included a term proportional to j_x and $\partial T_e / \partial x$, which was ascribed to a contribution from the Seebeck effect. Although the drift term derived above is similar in form to their term

$[\partial f_{\text{ex}} / \partial x = (\partial f_{\text{ex}} / \partial T_e)(\partial T_e / \partial x)]$, it has a different origin. The Seebeck effect vanishes in the present case of even-integer filling factors because the density of states is symmetric with respect to the chemical potential.

In stationary states, the equation for T_e becomes

$$M(\tilde{T}_e) \frac{d^2 \tilde{T}_e}{d\tilde{x}^2} = \eta(\tilde{T}_e) \frac{d\tilde{T}_e}{d\tilde{x}} + \tilde{T}_e - \gamma \tilde{j}_x^2 f_{\text{ex}}(1 - f_{\text{ex}}), \quad (19)$$

in a dimensionless unit where

$$M(\tilde{T}_e) = \kappa \hbar \omega_c / k_B C_L \ell^2, \quad (20)$$

$$\eta(\tilde{T}_e) = \eta_0 \tilde{j}_x f_{\text{ex}}(1 - f_{\text{ex}}) \tilde{T}_e^{-2}, \quad (21)$$

$$\eta_0 = eE_{c1} \omega_c / 2\pi \ell C_L, \quad (22)$$

$\tilde{x} = x/\ell$, $\tilde{j}_x = (-j_x)/j_{c1} = E_y/E_{c1}$, and $\gamma = 2.23342$. The estimation gives $\eta_0 = 50$ at $B = 5\text{ T}$. By changing the variable as $\tilde{t} = -\tilde{x}$, the above equation becomes the equation for “position” $\tilde{T}_e$ as a function of “time” $\tilde{t}$ of a particle with mass $M(\tilde{T}_e)$ in a position-dependent friction $\eta(\tilde{T}_e)$ and a potential $V(\tilde{T}_e)$ defined by

$$-\partial V / \partial \tilde{T}_e = \tilde{T}_e - \gamma \tilde{j}_x^2 f_{\text{ex}}(1 - f_{\text{ex}}), \quad (23)$$

and plotted in Fig. 1(a). We have chosen this direction of “time” because T_e with increasing $\tilde{x}$ approaches a stable value in a simple exponential form and is convenient for the initial point.

The current direction dependence is immediately obtained from the fact that $\eta \propto \tilde{j}_x$. The motion of the fictitious particle is much slower for the positive friction ($\tilde{j}_x > 0$) than for the negative friction ($\tilde{j}_x < 0$), reproducing the observed dependence of $T_e(x)$ on the current direction.

A critical behavior is expected to occur at $|j_x| = j_{c1}$, j_{c2} . We consider in this study the critical behavior at $|j_x| = j_{c1}$, since $j_{c2} = \infty$ in our simplified calculation of P_G and P_L . At $|j_x| = j_{c1}$, the homogeneous steady solutions are $\tilde{T}_e = 0$ and $\tilde{T}_e = \tilde{T}_{c1} = 0.324$. We consider below a relaxation from $T_e > T_{c1}$ ($x = 0$) to $T_e = 0$ ($x = \infty$) when $|j_x| < j_{c1}$ ($|\tilde{j}_x| < 1$).

First we show the critical behavior qualitatively using the above equivalent mechanical system. The initial “position” is chosen to be $\tilde{T}_{\text{init}} \equiv \tilde{T}_e(\tilde{x}_{\text{init}}) \ll \tilde{T}_{c1}$. At $\tilde{T}_e \sim \tilde{T}_{\text{init}}$, $f_{\text{ex}} \approx 0$ and $\tilde{T}_e = \tilde{T}_{\text{init}} \exp[-(\tilde{x} - \tilde{x}_{\text{init}})/M^{1/2}]$. Therefore the initial condition at $\tilde{x} = \tilde{x}_{\text{init}}$ is $\tilde{T}_e = \tilde{T}_{\text{init}}$ and $d\tilde{T}_e/d\tilde{t} = \tilde{T}_{\text{init}}/M^{1/2} > 0$, and the “particle” goes down the slope of the potential as shown in Fig. 1(a). When friction η is positive ($\tilde{j}_x > 0$), the motion of the particle slows down at the gradual slope at $\tilde{T}_e \sim \tilde{T}_{c1}$ and it takes a longer time to pass through the slope as $\tilde{j}_x \rightarrow 1$ since $\partial V / \partial \tilde{T}_e \rightarrow 0$ ($\tilde{j}_x \rightarrow 1$). This means the divergence of the relaxation length of $T_e(x)$ as $\tilde{j}_x \rightarrow 1$. On the other hand, no such critical behavior occurs when $\tilde{j}_x < 0$ since the friction is negative.

Next we calculate the length scale of the slow variation of $T_e(x)$. In this calculation, the second-derivative term can be neglected because the decay length l_d of $d^2 \tilde{T}_e / d\tilde{x}^2$ is given by $l_d = \ell M / \eta$ when $\partial V / \partial \tilde{T}_e = 0$ and is estimated to be $l_d = 5\text{ }\mu\text{m}$ at $B = 5\text{ T}$ and $T_e = T_{c1}$. The equation




Fig. 1. (a) “Potential” $V(\tilde{T}_e)$ defined in eq. (23) ($\tilde{T}_e = k_B T_e / \hbar \omega_c$, $\tilde{j}_x = (-j_x)/j_{c1} = E_y/E_{c1}$). (b) $\tilde{T}_e$ as a function of $\tilde{x} = x/\ell$ with η_0 defined in eq. (22) (thicker lines: numerical solutions of eq. (24), thinner lines: approximate analytical solutions, eq. (26)). (c) Relaxation length L_s defined in eq. (27) as well as h_0 , h_2 , and $\tilde{T}_0$ defined in eq. (25).

is integrated to give

$$\frac{\tilde{x} - \tilde{x}_0}{\eta_0} = -\tilde{j}_x \int_{\tilde{T}_0}^{\tilde{T}_e} d\tilde{T} \frac{1}{h(\tilde{T}, \tilde{j}_x)}, \quad (24)$$

with $h(\tilde{T}, \tilde{j}_x) = \tilde{T}^3(e^{1/2\tilde{T}} + e^{-1/2\tilde{T}} + 2) - \gamma \tilde{j}_x^2 \tilde{T}^2$. $\tilde{T}_0 = \tilde{T}_e(\tilde{x}_0)$ is the temperature at which $h(\tilde{T}, \tilde{j}_x)$ takes its minimum. The numerical integration gives $\tilde{T}_e(\tilde{x})$ plotted by thicker lines in Fig. 1(b), which shows that the length of the slow-variation region L_s increases as $\tilde{j}_x \rightarrow 1$. To find an analytical formula for L_s , we replace $h(\tilde{T}, \tilde{j}_x)$ by its Taylor series with respect to $\tilde{T}$ around its minimum up to the second order:

$$h(\tilde{T}, \tilde{j}_x) \approx h_0(\tilde{j}_x) + h_2(\tilde{j}_x)[\tilde{T} - \tilde{T}_0(\tilde{j}_x)]^2, \quad (25)$$

with $h_0(\tilde{j}_x)$, $h_2(\tilde{j}_x)$, and $\tilde{T}_0(\tilde{j}_x)$ plotted in Fig. 1(c). The analytical integration gives

$$\tilde{T}_e - \tilde{T}_0 = -\left(\frac{h_0}{h_2}\right)^{1/2} \tan \left[\frac{(h_0 h_2)^{1/2}}{\eta_0 \tilde{j}_x} (\tilde{x} - \tilde{x}_0) \right], \quad (26)$$

which is plotted in Fig. 1(b) by thinner lines; the curve

is in good agreement with the numerical ones. The relaxation length L_s is then defined by

$$L_s/\ell = \pi \eta_0 \tilde{j}_x (h_0 h_2)^{-1/2}. \quad (27)$$

L_s diverges as $(1 - \tilde{j}_x)^{-1/2}$ as shown in Fig. 1(c), and becomes $\sim 10 \mu\text{m}$ at $\tilde{j}_x = 0.97$.

The observed diagonal resistivity ρ_{xx} ¹¹⁾ remains to take large values within distance L_s^{exp} of approximately $50 \mu\text{m}$ ($\tilde{j}_x \approx 0.97$ and $B = 4.8 \text{ T}$) from the position of the hot-electron injection, and then undergoes a steep decrease. The calculated relaxation length, L_s , of T_e (corresponding to ρ_{xx}) is in agreement with the observed relaxation length, L_s^{exp} , in the order of magnitude and the critical behavior. The present theory under the assumption of $T_e \sim T_{c1}$ is not applicable to the observed steep decrease in the lower- ρ_{xx} (lower- T_e) region.

In conclusion, we have derived the hydrodynamic equation for the electron temperature T_e from the Boltzmann-type equation. We have applied the obtained equation to the spatial evolution of T_e and have reproduced two observed results: the current direction dependence and the critical behavior of the relaxation length for T_e .

The author would like to thank Y. Levinson, N. Tokuda, and Y. Asano for valuable discussions.

-
- [1] K. von Klitzing, G. Dorda and M. Pepper: Phys. Rev. Lett. **45** (1980) 494.
 - [2] S. Kawaji and J. Wakabayashi: in *Physics in High Magnetic Fields* (Springer, Berlin, 1981) p. 284.
 - [3] G. Ebert, K. von Klitzing, K. Ploog and G. Weimann: J. Phys. C **16** (1983) 5441.
 - [4] M. E. Cage, R. F. Dziuba, B. F. Field, E. R. Williams, S. M. Girvin, A. C. Gossard, D. C. Tsui and R. J. Wagner: Phys. Rev. Lett. **51** (1983) 1374.
 - [5] F. Kuchar, G. Bauer, G. Weimann and H. Burkhard: Surf. Sci. **142** (1984) 196.
 - [6] S. Kawaji, K. Hirakawa, M. Nagata, T. Okamoto, T. Fukase and T. Gotoh: J. Phys. Soc. Jpn. **63** (1994) 2303.
 - [7] G. Nachtwei: Physica E **4** (1999) 79.
 - [8] S. Komiyama and Y. Kawaguchi: Phys. Rev. B **61** (2000) 2014.
 - [9] S. Komiyama, Y. Kawaguchi, T. Osada and Y. Shiraki: Phys. Rev. Lett. **77** (1996) 558.
 - [10] Y. Kawaguchi: Doctor Thesis (University of Tokyo, 1997).
 - [11] I. I. Kaya, G. Nachtwei, K. von Klitzing and K. Eberl: Phys. Rev. B **58** (1998) R7536.
 - [12] I. I. Kaya, G. Nachtwei, K. von Klitzing and K. Eberl: Europhys. Lett. **46** (1999) 62.
 - [13] F. J. Ahlers, G. Hein, H. Scherer, L. Bliok, H. Nickel, R. Lösch and W. Schlapp: Semicond. Sci. Technol. **8** (1993) 2062.
 - [14] A. V. Gurevich and R. G. Mints: Pis'ma Zh. Eksp. Teor. Fiz. **39** (1984) 318 [JETP Lett. **39** (1984) 381].
 - [15] S. Komiyama, T. Takamasu, S. Hiyamizu and S. Sasa: Solid State Commun. **54** (1985) 479.
 - [16] H. Akera: J. Phys. Soc. Jpn. **69** (2000) 3174.
 - [17] J. A. Nixon and J. H. Davies: Phys. Rev. B **41** (1990) 7929.
 - [18] U. Wulf, V. Gudmundsson and R. R. Gerhardts: Phys. Rev. B **38** (1988) 4218.
 - [19] A. L. Efros: Solid State Commun. **67** (1988) 1019.
 - [20] A. H. MacDonald, T. M. Rice and W. F. Brinkman: Phys. Rev. B **28** (1983) 3648.
 - [21] S. Kawaji, K. Hirakawa and M. Nagata: Physica B **184** (1993) 17.
 - [22] A. Boisen, P. Bøggild, A. Kristensen and P. E. Lindelof: Phys. Rev. B **50** (1994) 1957.




Fig. 1 HiroshiAKE RA